

**Dr. med. Daniel A. Müller**

## Wann muss eine Weichteilschwellung abgeklärt werden?

---

### Bibliografie

- <sup>1</sup> O'Sullivan B, et al.: Preoperative versus postoperative radiotherapy in soft-tissue sarcoma of the limbs: a randomised trial. *Lancet* 2002; 359(9325): 2235–2241.
- <sup>2</sup> Collaboration SM-a: Adjuvant chemotherapy for localised resectable soft tissue sarcoma in adults. Sarcoma Meta-analysis Collaboration (SMAC). *Cochrane Database Syst Rev* 2000; 2.
- <sup>3</sup> Pretell-Mazzini J, et al.: Unplanned excision of soft-tissue sarcomas: current concepts for management and prognosis. *J Bone Joint Surg Am* 2015; 97(7): 597–603.

- <sup>4</sup> Jones DA, et al.: Management of Unplanned Excision for Soft-Tissue Sarcoma with Preoperative Radiotherapy Followed by Definitive Resection. *Am J Clin Oncol* 2016; 39(6): 586–592.
- <sup>5</sup> Chandrasekar CR, et al.: The effect of an unplanned excision of a soft-tissue sarcoma on prognosis. *J Bone Joint Surg Br* 2008; 90(2): 203–208.
- <sup>6</sup> Noria S, et al.: Residual disease following unplanned excision of soft-tissue sarcoma of an extremity. *J Bone Joint Surg Am* 1996; 78(5): 650–655.
- <sup>7</sup> Potter BK, et al.: Local recurrence of disease after unplanned excisions of high-grade soft tissue sarcomas. *Clin Orthop Relat Res* 2008; 466(12): 3093–3100.